

Wastewater as a Resource for Pest Control: An Overview

Ana M. Bailón-Salas,^{a,δ} Luis A. Ordaz-Díaz,^b Pablito M. López-Serrano,^a Monica Y. Flores-Villegas,^b and Pedro A. Dominguez-Calleros^{a,*}

Pests have a negative impact on the economy and the environment. There is an increased urgency for adequate pest control because many pests show high adaptation and climate change has created favorable circumstances for pests. For pest control, synthetic chemicals are used that are lethal to non-target organisms and are toxic to pollinators and aquatic invertebrates. Chemical compounds in plants and derivatives from lignocellulosic materials act against pests. The wastewater from lignocellulosic biomass is a potential source of new compounds with bactericidal, fungicidal, and pesticidal effects that have demonstrated inhibitory activity against plant pathogens. Fungicidal, nematocidal, insecticidal, larvicidal, and bactericidal activities have been proven. Inorganic and organic compounds, such as phenols, aldehydes, esters, and furanics, are the main ones identified. Due to the antimicrobial activity of wastewater, applying it to the soil can modify the composition and structure of key microbial communities. Deep research about richness, biodiversity, functionality, and microbials is needed. This review provides a comprehensive overview of wastewater types that have been applied and possible sources to obtain potential compounds for pest control. Moreover, associated active compounds, recovery techniques, and environmental impacts are reviewed.

Keywords: Biological control; Metabolites sources; Liquid waste; Lignocellulosic biomass

Contact information: a: Facultad de Ciencias Forestales, Doctorado Institucional en Ciencias Agropecuarias y Forestales, Río Papaloapan, Valle del Sur, C.P. 34120 Durango, Durango, México; b: Ingeniería en Tecnología Ambiental, Universidad Politécnica de Durango, Carr. Dgo-Mex Km 9.5, Col. Dolores Hidalgo, 34300, Durango, Durango, México; ^δ postdoctoral student; * Corresponding author: pdomingc@hotmail.com

CONTENTS

Introduction	2
Wastewater types for pest control	3
Cassava wastewater	3
Olive mill wastewater.....	3
Vinasses	5
Active compounds	5
Techniques for target compounds recovery	8
Environmental impacts	9
Positive impacts	10
Negative impacts.....	10
Conclusions	11

INTRODUCTION

Pests generate negative impacts because they decrease the quality and yield of crops (Savary *et al.* 2019). There is increased urgency because pests generate economic losses five times more than fires (Logan *et al.* 2003). Additionally, pathogens and pests are highly adaptable (Wingfield *et al.* 2015) and climate change can favor outbreaks and their extension (Rubin-Aguirre *et al.* 2015; Jactel *et al.* 2019). More so, the level of economic loss due to injuries caused by insects and pests is multifactorial, because they depend on the type of crop, temporal nature, and spatial location (Capinera 2020).

Plants contain chemical compounds that help fight disease and insects, for instance, phytohormones and secondary metabolites. The development of insecticides started from this discovery, and produced similar but more effective chemicals (Bednarek 2012; Erb *et al.* 2012; Matthews 2018). In contrast, lignocellulosic biomass is a copious and cheap source for pulp and paper, textile manufacturing, and agriculture in the forms of corn, wheat, rice, sorghum, barley, and sugarcane byproducts (Reddy and Yang 2005). Cellulose, hemicelluloses, and lignin are the main polymeric components (Sarip *et al.* 2016), the latter being the second most abundant biopolymer after cellulose (Demuner *et al.* 2019). In the lignocellulosic biomass pretreatment, the cellulose-hemicellulose-lignin structures are altered, facilitating the hydrolysis of cellulose and increasing the fermentable glucose concentration, whereby lignin derivatives are obtained (Kim 2018). Lignin contains several functional groups, such as phenolic hydroxyl, carboxylic, carbonyl, and methoxyl groups. Biological activities of phenolic hydroxyl and methoxyl groups (Espinoza-Acosta *et al.* 2016) were highlighted as antioxidant or antimicrobial activities (Alzagameem *et al.* 2019; Jinxiang *et al.* 2020). In addition, chemical compounds derived from the by-products from lignocellulosic materials provide protection for pests (Villaverde *et al.* 2016).

For pest control, methyl bromide gas was widely used as a broad-spectrum fumigant until 2005, when it was banned (Wedge *et al.* 2001). Neonicotinoid pesticides have become the most widely used class of insecticides in the world (Simon-Delso *et al.* 2014). However, they present significant environmental impacts (Saeed *et al.* 2019). For example, methyl bromide produces neurotoxicity and is a stratospheric ozone depleter (Wedge *et al.* 2001), the fipronil is toxic to pollinators and aquatic invertebrates (Sadaria *et al.* 2019), and most insecticides can be lethal to non-target organisms (Simon-Delso *et al.* 2014). In addition, pesticides are considered a powerful biological risk because they can persist in the environment for years (Sharma *et al.* 2020). Chemical fungicides are widely used because they are effective in sterilizing (Lin *et al.* 2020). However, they have also been reported to induce resistance in fungal plant pathogens (Swett *et al.* 2020). For the control of phytoparasitic nematodes, fumigants have been used that have the ability to eliminate not only target organisms but also affect the microbial population in the soil (Ntalli *et al.* 2020). This leads researchers to search for alternatives to control pests.

The use of living biological organisms or their metabolites for pest control is called bio-pesticides (Butu *et al.* 2020). The demand is increasing to limit the use of chemical pesticides and to replace them with agents that have no or less negative effects on the environment (Di Ilio and Cristofaro 2020; Rashwan and Hammad 2020). Sharma *et al.* (2020) considered that biopesticides could facilitate increased crop production with or without minimal negative effects. Furthermore, biopesticides are biodegradable, less expensive, and possess less toxicity toward living organisms (Thakur *et al.* 2020).

There are different alternatives for pest control, such as genetically based resistance (Molinari 2011), integrated pest management (IPM) (Meissle *et al.* 2010), botanical

pesticides (Lengai *et al.* 2020), larvicidal red-algae (Deepak *et al.* 2019), reductive soil disinfestation (RSD), anaerobic soil disinfestation (Huang *et al.* 2019), and microbial biopesticides (Thakur *et al.* 2020), among others. Biopesticide sources exist readily in nature (Thakur *et al.* 2020), and some have yet to be fully exploited and studied, for example, liquid waste.

Wastewater is a valuable source of biomolecules for different uses; by recovering these compounds, value is added, and at the same time the environmental impact in the treatment of these wastes is reduced (Larif *et al.* 2015; Ahmad *et al.* 2020). These wastes may possibly be considered one of the most abundant, cheap, and renewable resources on earth (Gonzalez-Coloma *et al.* 2013). It is time to change the paradigm and stop seeing them only as waste to treat them as by-products, revalue the “waste” and give them another type of value with sustainable management of these materials (Ordaz-Díaz *et al.* 2019).

Wastewater Types for Pest Control

The raw wastewaters used for pest control are cassava, olive mill, vinasses of wine, sugar beet, and sugarcane featuring fungicidal, nematicidal, insecticidal, larvicidal, and bactericidal activities (Table 1).

Cassava Wastewater

A feature of cassava wastewater is the presence of linamarin and lotaustralin compounds, cyanogenic glycosides that are lost in processing (Padmaja 1995). Near 5 to 7 L of wastewater are generated from a kilogram of fresh cassava root (Wathier *et al.* 2019). The cassava wastewater has been studied for more than 30 years for possible applications in pest management (Pinto-Zevallos *et al.* 2018), against insects, nematodes, and fungi (Table 1). The pest species that have been evaluated are *Coceus hesperidum* L., *Meloidogyne* spp., and *Oidium* sp., which are associated mainly with crops of fruit trees and tomato (Lebeda *et al.* 2015; Abdul-Rassoul 2016; Regmi and Desaegeer 2020).

Olive Mill Wastewater

Olive mill wastewater contains phenolic compounds (Di Mauro *et al.* 2017). Due to the reducing power of these compounds, bacteria and plants are negatively impacted (Babić *et al.* 2019). The olive mill wastewater can be used against bacterial, fungal phytopathogens, and weed species (El-Abbassi *et al.* 2017). Pest mortality is attributed to phenolic compounds (Larif *et al.* 2013). Hence, polyphenolic fractions of the olive mill wastewater act as a strong natural chemosterilant (Di Ilio and Cristofaro 2020).

Euphyllura olivina and *Ceratitis capitata* Wiedemann, globally important pests, are Mediterranean parasitoids that affect olive and fruit crops, respectively (Alves *et al.* 2019; Hougardy *et al.* 2020). The olive mill wastewater has also shown insecticidal activity against both pests and larvicidal activity against *Euphyllura olivina*. Furthermore, *Aphis citricola*, an aphid related to apple orchards infestation (Kou *et al.* 2020), can be controlled using olive mill wastewater, due to larvicidal activity (Table 1).

Besides being effective against larvae and insects, the olive mill wastewater has also been shown to suppress fungi and bacteria activity (Table 1).

Table 1. Wastewater Type Used for Pest Control

Pest	Activity	Wastewater Type	Reference
<i>Phytophthora parasitica</i> , <i>Fusarium oxysporum</i> f. sp. <i>melonis</i> race, <i>F. oxysporum</i> f. sp. <i>radicis-cucumerinum</i> , <i>Pythium aphanidermatum</i> , and <i>Sclerotinia sclerotiorum</i>	Fungicidal	Sugar beet, sugarcane, and wine vinasse	Santos <i>et al.</i> 2008
<i>Meloidogyne incognita</i>	Nematicidal	Sugar beet vinasse	Núñez-Zofío <i>et al.</i> 2013
<i>Sphenophorus levis</i>	Insecticidal	Sugarcane vinasse	Martins <i>et al.</i> 2020
<i>Stomoxys calcitrans</i>			Jelvez Serra <i>et al.</i> 2017
<i>Oregmomyga peruviana</i>	Insecticidal	Wine vinasse	Dadther-Huaman <i>et al.</i> 2020
<i>Coccus hesperidum</i> L.	Insecticidal	Cassava	Ponte <i>et al.</i> 1988
<i>Meloidogyne</i> spp.	Nematicidal		Ponte and Franco 1983
<i>Oidium</i> sp.	Fungicidal		Santos and Ponte 1993
<i>Euphyllura olivina</i>	Insecticidal	Olive mill	Debo <i>et al.</i> 2011
<i>Ceratitis capitata</i> Wiedemann			Di Ilio and Cristofaro 2020
<i>Euphyllura olivina</i> and <i>Aphis</i> <i>citricola</i>	Larvicidal		Larif <i>et al.</i> 2013
<i>Botrytis cinerea</i>	Fungicidal		Yangui <i>et al.</i> 2010
<i>Rhizoctonia solani</i> and <i>Fusarium oxysporum</i>			Mohamed <i>et al.</i> 2015
<i>Fusarium sambucinum</i> , <i>Verticillium</i> <i>dahliae</i> , and <i>Alternaria</i> <i>solani</i>			Yangui <i>et al.</i> 2009
<i>Pseudomonas syringae</i> and <i>Xanthomonas campestris</i>			Bactericidal

The fungicidal activity of olive mill wastewater has been tested against *Botrytis cinerea*, *Rhizoctonia solani*, *Fusarium oxysporum*, *Fusarium sambucinum*, *Verticillium dahlia*, and *Alternaria solani* (Table 1). These fungi affect various crops and the economic losses they generate are considerable. For example, *Botrytis cinerea*, a necrotrophic pathogen, produces severe crop losses worldwide in a wide variety of plant species (Hahn *et al.* 2014). *Rhizoctonia solani* is a root pathogen that affects cereal crops (Paulitz and Schroeder 2005). *Fusarium oxysporum* is a soil and seed-borne disease and is one of the main pathogens of dry rot (Tiwari *et al.* 2020). *Fusarium sambucinum* (root rot disease) and *Fusarium oxysporum* cause potato infection (Yangui *et al.* 2009; Pilsyk *et al.* 2015; Tiwari *et al.* 2020). *Verticillium dahliae* is a vascular pathogen that causes wilt and death of 400 cultivated and non-cultivated plant species including the tomato plants. *Alternaria solani* affects different parts of the plant from root rot to even cause tomato and potato rot (Yangui *et al.* 2009; EFSA Panel on Plant Health PLH 2014).

Pseudomonas syringae is an extracellular bacteria and is considered one of the main bacterial pathogens of plants (Mansfield *et al.* 2012; Xin *et al.* 2018). *Xanthomonas campestris* is a bacteria able to cause black rot infection in cruciferous plants (Papaianni *et al.* 2020). Due to the bactericidal activity of olive mill wastewater, both phytopathogenic bacteria are inhibited by this liquid waste (Table 1).

Vinasses

The vinasse, a liquid residue from alcoholic fermentation, contains various compounds such as alcohols, aldehydes, phenols, and acids (Couallier *et al.* 2006; Freitas *et al.* 2018; Fuess *et al.* 2018). In some cases, these compounds are undesirable. For example, in anaerobic wastewater treatment, phenolic compounds should be removed, because they participate as inhibitors (Borja *et al.* 1993; Ao *et al.* 2020). However, some of the compounds have a positive environmental and economic value.

Raw vinasse has proven useful in other fields of research. Phanapavudhikul (1999) observed an eradication of insects by adding the sugarcane vinasse to the soil, associating it with oxygen depletion. The first reports of vinasse use to control phytopathogenic fungi date back to 2008. It was reported that wine vinasse showed 100% efficacy in suppressing the growth of phytopathogenic fungi (Santos *et al.* 2008). Afterward, the sugar beet vinasse was tested for the control of nematodes in pepper crops, as an alternative to the disinfection of soil-borne pathogens (Núñez-Zofío *et al.* 2013). Furthermore, the vinasse compounds can be used as chemical attractants (Martins *et al.* 2020). Recently, in the treatment of mycoremediation, Fernandes *et al.* (2020) reported a decrease in the growth rate of fungi using a wine vinasse concentration higher than 60%.

The vinasses have been shown to be effective against fungi, insects, and nematodes. The fungicidal activity have been tested against *Phytophthora parasitica*, *Fusarium oxysporum* f. sp. *melonis* race, *F. oxysporum* f. sp. *radicis-cucumerinum*, *Pythium aphanidermatum*, and *Sclerotinia sclerotiorum* (Table 1). The *Phytophthora* genus is one of the most devastating pathogens to a wide range of crop plants (El-Sayed and Ali 2020). *Phytophthora parasitica* is a soilborne pathogen (Meng *et al.* 2014). This oomycete mainly affects tobacco (Hou *et al.* 2012), tomato crops (Vigo *et al.* 2000), and the citrus industry (Boava *et al.* 2011). *Fusarium oxysporum* f. sp. *melonis* race is one of the most important diseases causing tremendous losses in melon fruit (Almasi 2019). *F. oxysporum* f. sp. *radicis-cucumerinum* is a vascular wilt fungus and is associated with cucumber crops (Markakis *et al.* 2016). *Pythium aphanidermatum* is the most devastating pathogen that affects turmeric and *Sclerotinia sclerotiorum* is capable of attacking more than 400 crop species (Boland and Hall 1994; Chand *et al.* 2016).

Moreover, the insecticidal activity was evaluated using sugarcane and wine vinasse against *Sphenophorus levis*, *Stomoxys calcitrans*, and *Oregmomyza peruviana* (Table 1). *Sphenophorus levis* affects the sugarcane crops, *Stomoxys calcitrans* is a stable fly that acts as a mechanical vector for the lumpy skin disease virus on cattle, and *Oregmomyza peruviana* is a vine pest (Casteliani *et al.* 2020; Dadther-Huaman *et al.* 2020; Paslaru *et al.* 2020).

Meloidogyne incognita, a root-knot nematode damaging vegetable crops (Collange *et al.* 2011), also has been tested using sugar beet vinasse for pest control (Table 1).

Due to this potential, vinasse can be studied as a source of biocide for the prevention and control of various pests.

Active Compounds

Esters, acids, aldehydes, ketones, aromatics, alkanes, alcohols, nitrosamides, and terpenoids, are acting in a synergistic inhibitory manner of fungi and bacteria (Saxena and Strobel 2020). Table 2 shows the compounds present in wastewater, a diverse source (vinasses, olive mill, and cassava) that is of great interest for the control pests. Phenols, organic acids, aldehydes, esters, furanic, and inorganic compounds are the main ones.

Table 2. Compounds Identified in Wastewater Samples of Interest in Pest Control

Wastewater Type	Classification	Compounds	Content	Reference	
Wine Vinasses	Phenol	Gallic acid	10.83 g /L	Díaz <i>et al.</i> 2012	
		Hydroxytyrosol	ND		
		Total phenols	18.9 g/L		
Baijiu Vinasse			Ferulic acids	1.674 mg/K	Wang <i>et al.</i> 2019
Sugar Beet Vinasse			Ferulic acid	< 5 mg/L	Bostyn <i>et al.</i> 2009
Tequila Vinasse			Eugenol	0.9 mg/L	Félix <i>et al.</i> 2018
			2,4-di-tert-butylphenol	90 mg/L	
			4-(2-hydroxyethyl) phenol	ND	
Mezcal Vinasses			Gallic acid	478 to 542 mg/L	Robles-González <i>et al.</i> 2012
Olive Mill Wastewater			Gallic acid	ND	Puoci <i>et al.</i> 2012
		Oleuropein	14.32		
			Hydroxytyrosol	267.17 to 821.86 mg/L	Di Mauro <i>et al.</i> 2017
Tequila Vinasse	Aldehydes	Benzaldehyde	ND	Félix <i>et al.</i> 2018	
	Esters	Ethyl butanoate, ethyl lactate, and ethyl palmitate	ND		
	Alkanes	Dodecane, tetradecane, and eicosane	ND		
	Furanic compounds	Furfural	50 mg/L		
		5-methyl furfural	ND		
Pyrans	4H-pyran-4-one,2,3-dihydro-3,5-dihydroxy-6-methyl and pyrrolo[1,2-a] pyrazine-1,4-dione, hexahydro-3-(2-methylpropyl)	ND			
Sugarcane Vinasse	Organic acids	Lactic acid	1.2X10 ⁻¹ mol/L	Sedenho <i>et al.</i> 2017	
Cassava Wastewater	Triazine	Cyanuric acid	ND	Pinto-Zevallos <i>et al.</i> 2018	
		Free cyanide	257.20 mg/L	Neves <i>et al.</i> 2014	
Sugarcane Vinasse	-	Melanoidins	16600 g/L	Kaushik <i>et al.</i> 2018	

ND=no data

Pest mortality has been attributed to the presence of phenolic compounds (Larif *et al.* 2013), due to being part of the protection system of plants against pests (Patzke and Schieber 2018). Lignin or lignin-rich biomass are a source of phenols, which can be obtained through the hydrothermal process (Peng *et al.* 2019). Therefore processes that contain lignin and are subjected to high temperatures will contain phenolic compounds in their wastewater, thanks to thermal hydrolysis. The wastewater contains phenols, such as hydroxytyrosol, gallic acid, ferulic acid, eugenol, oleuropein, 2,4-di-tert-butylphenol, and 4-(2-hydroxyethyl) phenol, which can be used or recovered for pest control (Table 2). For instance, hydroxytyrosol metabolite, a phytochemical polyphenol with antioxidant

properties, has exhibited antimicrobial activity (Bisignano *et al.* 1999), insecticidal activity (Debo *et al.* 2011), disinfectant activity on seeds (Yangui *et al.* 2009), and fungicide activity (Yangui *et al.* 2010; Khan and Murphy 2020). The main sources of hydroxytyrosol are olive and wine (Rebollo-Romero *et al.* 2020). However, it is also reported in wine vinasse and olive mill wastewater (Table 2). Gallic acid, a secondary polyphenolic metabolite, is considered one of the most powerful antioxidants and has been reported in most plants (Erukainure *et al.* 2018; Martínez *et al.* 2018). From an environmental point of view, gallic acid present in agro-industrial wastewaters must be removed, due to its toxicity (V́ctor-Ortega and Airado-Rodŕguez 2018), because it can affect the microbial communities in wastewater discharge points. However, it can be used for the management of bacteria pathogen pests, as Borges *et al.* (2013) reported bactericidal activity. This phenolic compound is available in olive mill wastewater, wine, and mezcal vinasses, as an alternate source (Table 2). Ferulic acid is a phenolic compound extremely abundant and found widely in nature (Rosazza *et al.* 1995), showing fungicidal and bactericidal activities (Borges *et al.* 2013; Patzke and Schieber 2018). Based on Table 2, this compound has been reported in baijiu and sugar-beet vinasses. The eugenol (4-allyl-2-methoxyphenol) found in tequila vinasse (Table 2); it is an acaricidal agent, having fungicidal and bactericidal activities (Abd El-Baky and Hashem 2016; Shang *et al.* 2020). Oleuropein shows bactericidal and fungicidal activities, mainly contained in olives (Bisignano *et al.* 1999), and, as shown in Table 2, was also in the wastewater. Additionally, 2, 4-di-tert-butylphenol demonstrated fungicidal and acaricidal activity (Dharni *et al.* 2014; Chen and Dai 2015; Varsha *et al.* 2015), and 4-(2-hydroxyethyl) phenol showed nematocidal activity (Yang *et al.* 2012).

Therefore, phenolic compounds are considered a natural alternative to conventional plant protection agents (Patzke and Schieber 2018). Additionally, the recovery of phenols and the obtaining of added value products is attractive from industrial and environmental points of view (V́ctor-Ortega and Airado-Rodŕguez 2018).

Tequila vinasse compounds such as aldehydes, esters, alkanes, furanic compounds, and pyrans can be used for pest control. The benzaldehyde identified in tequila vinasse (Table 2) could be applicable for developing novel insecticides for agricultural use due to having been tested as agents effectively inhibiting fungi, insects, and microbials (Kim *et al.* 2011; Ullah *et al.* 2015). This compound is present in *Agave* alcoholic beverages such as bacanora, mezcal from *A. angustifolia*, mezcal from *A. durangensis*, mezcal from *A. potatorum*, mezcal from *A. salmiana*, raicilla, sisal, sotol, tequila, and pulque (De León Rodríguez *et al.* 2008). Therefore it is also found in the vinasse of each process. The ethyl butanoate, an ester, was identified in male rectal glands during periods of sexual activity in the banana fruit fly (*Bactrocera musae* Tryon). Therefore, ethyl butanoate could be used to control this pest as a possible biological role of these compounds in the mating system (Noushini *et al.* 2020). The ethyl lactate can be generated from biomass raw materials through fermentation (Pereira *et al.* 2011). As shown in Table 2, it is also present in liquid distillation wastes of tequila. A mix of ethyl lactate and acetic acid exhibits an antifungal effect (Steven *et al.* 2016). Ethyl palmitate was identified as a component of the pheromone from the brood of bees, and this volatile compound attracts the small hive beetle, a pest of honeybees. This finding can be useful for trap development and management (Dekebo and Jung 2020). Other compounds present in the tequila vinasse, such as dodecane, tetradecane, and eicosane (Table 1), are the female sex pheromone compounds of *Paranthrene diaphana* Dalla Torre and Strand (Lep. Sesiidae), a destructive pest of willow trees (Minaeimoghadam *et al.* 2017). Therefore, the vinasse could be used to attract and control

this pest. Moreover, furanic compounds with antifungal activity, such as furfural and 5-methyl furfural, can help decrease the commercial antifungal agrochemical dose against *Alternaria mali* (Jung *et al.* 2007). Additionally, 4H-pyran-4-one, 2,3-dihydro-3,5-dihydroxy-6-methyl (Table 2) is part of a mixture that shows insecticidal, larvicidal, and pupicidal effects (Ravindran *et al.* 2020), and pyrrolo[1,2-a] pyrazine-1,4-dione, hexahydro-3-(2-methylpropyl) (Table 2) has an antifungal function (Kannabiran 2016).

Lactic acid identified in sugarcane vinasse (Table 2) showed antifungal activity against *Aspergillus*, *Penicillium*, and *Fusarium* genera (Lind *et al.* 2005). L-Lactic acid has used as a pesticide and can be obtained through a fermentation process (Liu *et al.* 2013). The cyanide that is present in cassava wastewater (Table 2) acts as a natural plant defense against pests (Pinto-Zevallos *et al.* 2018). This is the case for the cyanogenic glycosides in wastewater that comes from the soaking stage in the manufacture of flour from cassava (Alitubeera *et al.* 2019).

Melanoidins are the end products of the Maillard reaction between carbohydrates and amino compounds (Cämmerer and Kroh (1995). They are found in vinasses (Table 2), and have antimicrobial activity (Kaushik *et al.* 2018).

It can be seen in Table 2 that the composition of wastewater is diverse. Koul and Walia (2009) mentioned that this can be an advantage because the possibility of pests developing resistance is reduced.

Techniques for Target Compounds Recovery

The direct application of raw wastewater has been the most used for the evaluation of the power against pests. However, it is possible to recover the target compounds.

For the compound's recovery in rich phenolic wastewater, magnetic extraction, ultrasound-assisted extraction, solvent extraction, adsorption, or combined processes, such as hydrolysis-purification and extraction-adsorption, are used (Table 3). However, sometimes the suspended matter needs to be removed by flocculation, as a preliminary stage (Azzam and Hazaimh 2021).

Some of the solvent extraction process steps are acidification or condensing, delipidation extraction, and purification used for phenol recovery in olive mill wastewater (Deng *et al.* 2017; Çelik *et al.* 2020). The acidification with acetic acid allows hydroxytyrosol enrichment (Debo *et al.* 2011), and ultrasound-assisted extraction could increase the yield of phenolic compounds (Deng *et al.* 2017). In the ultrasound-assisted extraction, less solvent is required (Albero *et al.* 2015), which makes it a more environmentally friendly technique. To remove lipids, a fraction delipidation step is employed, and hexane is the most used (Rubio-Senent *et al.* 2017). Ethyl acetate is the solvent more commonly used for the recovery of high added-value compounds from wastewaters (Table 3). This solvent was the most efficient for the recovery of phenolic monomers from olive mill wastewater (Allouche *et al.* 2004), and the system was able to reach a total recovery of polyphenols (Bostyn *et al.* 2009). After extraction, the resins are used in the purification step to increase the amount and purity of phenolic compounds (Çelik *et al.* 2020).

Despite adsorbents or chemicals used in conventional treatment (absorption and extraction) are cheaper than advanced treatments, both show high efficiencies (Villegas *et al.* 2016). It is even possible to reduce costs further with the use of low-cost adsorption media (Daragon *et al.* 2014).

Table 3. Processes Used for the Recovery of Phenol in Wastewater

Wastewater Type	Compounds	Solvent	pH	Process	Reference
Olive Mill Wastewater	Hydroxytyrosol	Ethyl acetate	2	Solvent extraction	Allouche <i>et al.</i> 2004
		-	3	Hydrolysis (acetic acid) and purification (resin)	Debo <i>et al.</i> 2011
	Polyphenols	Ethyl acetate	-	60 °C	Larif <i>et al.</i> 2015
Tequila Vinasse	Phenolic compounds			Adsorption resins	Sanchez <i>et al.</i> 2019
Sugar Beet Vinasse	Phenolic acids	Ethyl acetate	4	Solvent extraction	Bostyn <i>et al.</i> 2009
Olive Oil Wastewater	Gallic acid	Ethanol		Solvent extraction and adsorption (molecularly imprinted polymers)	Puoci <i>et al.</i> 2012
Aqueous Solutions	Gallic acid	Ethyl acetate	-	Solvent extraction	Daneshfar <i>et al.</i> 2008
	Phenolic compounds	-	-	Magnetic extraction	Deng <i>et al.</i> 2011
Urban wastewater	Phenolic compounds	-	-	Ultrasound-assisted	Kotowska <i>et al.</i> 2014

Table 4 shows that certain wastewater types can have high lignin content, such as wheat straw and kraft pulping effluent. In fact, the dark color in cassava wastewater is due to the presence of lignin breakdown products and lignin phenols (Zhang *et al.* 2017). Lignin can be used as a raw material in the production of aromatic monomers (Gu *et al.* 2020). Currently, catalytic hydrothermal depolymerization has been used to obtain phenolic monomers from lignin (Roy *et al.* 2020), with the yield increasing 49% and 83% when using mannitol and sucrose addition (Gu *et al.* 2020).

Table 4. Lignocellulosic Biomass in Wastewater

Wastewater Type	Polymer	Content	Reference
Olive mill	Lignin	25.5 *	Uğurlu and Kula 2007
Wheat straw	Lignin	310 to 660 *	Wang 2020
Aspen kraft pulp	Lignin	230 to 770 *	
Sorghum and rice vinasse	Lignin	14.95 **	Cao <i>et al.</i> 2019

*mg/L, **% Dry matter

Environmental Impacts

For pest control, the use of compounds in wastewater should be discussed relative to potential environmental impacts. This source is a new field of research and the possible negative or positive effects have not been studied in depth.

As previously discussed, the positive effect of wastewater use on pests has been shown. However, when the wastewater is in contact with parts of the plants, with the soil,

or is infiltrated into underground water, the effects are unclear. Possible environmental negative and positive impacts of the use of wastewater for pest control are shown in Table 5.

Positive Impacts

The positive impact, in general, is that wastewaters rich in nutrients are considered as an alternative fertilizer, bringing enhanced crop growth, water-holding capacity improvement, and microbial communities in wastewater coadjutant to phenolic degradation in soil (Table 5). Moreover, better soil basal respiration has been reported in cassava wastewater (Table 5), and the increase in CO₂ produced in soil respiration is associated with the improvement biological activity of organisms (Phillips and Nickerson 2015). The treated olive mill wastewater germination index increases and water-holding capacity with raw wastewater is improved and does not contain high heavy metal levels (Table 5).

Table 5. Environmental Impacts of Wastewater Used as Pest Control

Wastewater Type	Environmental Impact		Reference
	Positive	Negative	
Cassava	Increased nutrients	Initial toxic effect on soil fungal	dos Santos Moura <i>et al.</i> 2018
	Total organic carbon increase		
	Microbial biomass increase		
	Better soil basal respiration		
	Soil fertility improvement	Soil hydrophobicity	Abegunrin <i>et al.</i> 2016
	Enhanced crop growth		
	Crop yield increase		Cabral <i>et al.</i> 2010
Olive Mill	Germination index increase		Mekki <i>et al.</i> 2013
	Water-holding capacity improvement		
	Low heavy metal loads	Seeds germination	Sassi <i>et al.</i> 2006
	Increased nutrients and organic matter	Soil infiltration rates decrease	Zema <i>et al.</i> 2019
		Soil infiltration rates decrease	Albalasmeh <i>et al.</i> 2019
		Microbial communities in olive mill wastewater coadjutant to phenolic degradation in soil	El Hassani <i>et al.</i> 2020
Sugar Beet Vinasse		Vegetation cover decreasing	Tejada <i>et al.</i> 2009
Sugarcane Vinasse		Methane emissions	Do Carmo <i>et al.</i> 2012
	Organic carbon increase		Soobadar and Ng Kee Kwong 2012
	Crop yield increase		

Negative Impacts

The raw and concentrated wastewater show negative impacts, for instance, concentrated olive mill wastewater blocked seed germination, and an initial toxic effect on soil fungal activity were identified (Table 5). Sassi *et al.* (2006) reported a dilution of 1/16 to guarantee full germination. High salt content in vinasse can cause vegetation cover decrease and changes in the structure and porosity of the soil (Tejada *et al.* 2009). When

wastewaters are added to the soil, the organic matter accumulation in the soil affects penetration resistance and water repellency (Albalasmeh *et al.* 2019). A pre-treatment is recommended before applying it to the soil (Abegunrin *et al.* 2016). To avoid soil hydrophobicity and methane emissions (Table 5), a flocculation process can help.

Wastewater reviewed in this paper contained compounds with antimicrobial activity (Table 2). Therefore, care must be taken when applying it to the soil, as it can modify the composition and structure of key microbial communities. Hence, more research addressed toward microbial communities' impacts on the soil is needed, with a focus on richness, biodiversity, functionality, and microbial adaptability.

To avoid most of the negative environmental impacts associated with raw and concentrated wastewater for pest control, compound recovery processes with fungicidal, acaricidal, nematicidal, bactericidal, and insecticidal activities can be exploited. However, it is necessary to assess the economic and environmental benefits of both options, with more in-depth research.

Otherwise, there is another wastewater type with compounds that can inhibit pests; for example, pulp and paper mill effluent, which is a chlorophenol source (Cheng *et al.* 2015). Chlorophenols used as herbicides or fungicides on crops were banned, due to human carcinogenic risk (Owuor 2003; Badanthadka and Mehendale 2014). Therefore, pulp and paper mill effluent is not susceptible to use as a source for control pests.

Humans and animals as part of the environment can also be affected by the use of compounds that help control pests and plant diseases. Lin *et al.* (2020) mention that the use of chemical fungicides in fruits or vegetables could modify the composition of the intestinal microbiota. Therefore, a more complete study is required on the application of wastewater in pest control, either as a direct application or compounds recovery of interest, considering the present economic-environmental impact.

CONCLUSIONS

1. Wastewater is a potential and alternative source of compounds with bactericidal, fungicidal, and pesticidal effects that have demonstrated inhibitory activity.
2. The phenolic compounds are mainly responsible for pest mortality using wastewater.
3. Because wastewater is a variable chemical composition, its use can be dangerous for the environment. Therefore, the isolation of target compound is recommended.

ACKNOWLEDGMENTS

The support of the Consejo Nacional de Ciencia y Tecnología (CONACyT) is gratefully appreciated for the support of the Sistema Nacional de Investigadores (SNI) and the postdoctoral scholarship (130780/20) received by one of the writers.

REFERENCES CITED

- Abd El-Baky, R. M., and Hashem, Z. S. (2016). "Eugenol and linalool: Comparison of their antibacterial and antifungal activities," *African Journal of Microbiology Research* 10(44), 1860-1872. DOI: 10.5897/AJMR2016.8283
- Abdul-Rassoul, M. S. (2016). "New host plants record for the brown soft scale *Coccus hesperidum* Linnaeus, 1758 (Hemiptera: Coccidae) in Baghdad province, Iraq," *Bulletin of the Iraq Natural History Museum* 14(2), 135-139.
- Abegunrin, T. P., Awe, G. O., Idowu, D. O., and Adejumobi, M. A. (2016). "Impact of wastewater irrigation on soil physico-chemical properties, growth and water use pattern of two indigenous vegetables in southwest Nigeria," *CATENA* 139, 167-178. DOI: 10.1016/j.catena.2015.12.014
- Ahmad, T., Belwal, T., Li, L., Ramola, S., Aadil, R. M., Xu, Y., and Zisheng, L. (2020). "Utilization of wastewater from edible oil industry, turning waste into valuable products: A review," *Trends in Food Science & Technology* 99, 21-33. DOI: 10.1016/j.tifs.2020.02.017
- Albalasmeh, A. A., Alajlouni, M. A., Ghariabeh, M. A., and Rusan, M. J. (2019). "Short-term effects of olive mill wastewater land spreading on soil physical and hydraulic properties," *Water, Air, & Soil Pollution* 230(8), article no. 208. DOI: 10.1007/s11270-019-4243-5
- Albero, B., Sánchez-Brunete, C., García-Valcárcel, A. I., Pérez, R. A., and Tadeo, J. L. (2015). "Ultrasound-assisted extraction of emerging contaminants from environmental samples," *TrAC Trends in Analytical Chemistry* 71, 110-118. DOI: 10.1016/j.trac.2015.03.015
- Alitubeera, P. H., Eyu, P., Kwesiga, B., Ario, A. R., and Zhu, B. P. (2019). "Outbreak of cyanide poisoning caused by consumption of cassava flour—Kasese District, Uganda, September 2017," *Morbidity and Mortality Weekly Report* 68(13), 308. DOI: 10.15585/mmwr.mm6813a3
- Allouche, N., Fki, I., and Sayadi, S. (2004). "Toward a high yield recovery of antioxidants and purified hydroxytyrosol from olive mill wastewaters," *Journal of Agricultural and Food Chemistry* 52(2), 267-273. DOI: 10.1021/jf034944u
- Almasi, M. A. (2019). "Development of a colorimetric loop-mediated isothermal amplification assay for the visual detection of *Fusarium oxysporum* f. sp. *melonis*," *Horticultural Plant Journal* 5(3), 129-136. DOI: /10.1016/j.hpj.2019.01.004
- Alves, T. J. S., Murcia, A., Wanumen, A. C., Wanderley-Teixeira, V., Teixeira, Á. A. C., Ortiz, A., and Medina, P. (2019). "Composition and toxicity of a mixture of essential oils against Mediterranean fruit fly, *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae)," *Journal of Economic Entomology* 112(1), 164-172. DOI: 10.1093/jee/toy275
- Alzagameem, A., Klein, S. E., Bergs, M., Do, X. T., Korte, I., Dohlen, S., Hüwe, C., Kreyenschmidt, J., Kamm, B., Larkins, M., et al. (2019). "Antimicrobial activity of lignin and lignin-derived cellulose and chitosan composites against selected pathogenic and spoilage microorganisms," *Polymers* 11(4), article no. 670. DOI: 10.3390/polym11040670
- Ao, T., Luo, Y., Chen, Y., Cao, Q., Liu, X., and Li, D. (2020). "Towards zero waste: A valorization route of washing separation and liquid hot water consecutive pretreatment to achieve solid vinasse based biorefinery," *Journal of Cleaner Production* 248, article ID 119253. DOI: 10.1016/j.jclepro.2019.119253

- Azzam, M. O. J., and Hazaimh, S. A. (2021). "Olive mill wastewater treatment and valorization by extraction/concentration of hydroxytyrosol and other natural phenols," *Process Safety and Environmental Protection* 148, 495-523. DOI: 10.1016/j.psep.2020.10.030
- Babić, S., Malev, O., Pflieger, M., Lebedev, A. T., Mazur, D. M., Kužić, A., Čož-Rakovac, R., and Trebše, P. (2019). "Toxicity evaluation of olive oil mill wastewater and its polar fraction using multiple whole-organism bioassays," *Science of The Total Environment* 686, 903-914. DOI: 10.1016/j.scitotenv.2019.06.046
- Badanthadka, M., and Mehendale, H. M. (2014). "Chlorophenols," in: *Encyclopedia of Toxicology*, P. Wexler (ed.), Elsevier Academic Press, Amsterdam, Netherlands, pp. 896-899. DOI: 10.1016/B978-0-12-386454-3.00281-5
- Bednarek, P. (2012). "Chemical warfare or modulators of defense responses – The function of secondary metabolites in plant immunity," *Current Opinion in Plant Biology* 15(4), 407-414. DOI: 10.1016/j.pbi.2012.03.002
- Bisignano, G., Tomaino, A., Cascio, R. L., Crisafi, G., Uccella, N., and Saija, A. (1999). "On the *in-vitro* antimicrobial activity of oleuropein and hydroxytyrosol," *Journal of Pharmacy and Pharmacology* 51(8), 971-974. DOI: 10.1211/0022357991773258
- Boava, L. P., Cristofani-Yaly, M., Mafra, V. S., Kubo, K., Kishi, L. T., Takita, M. A., Ribeiro-Alves, M., and Machado, M. A. (2011). "Global gene expression of *Poncirus trifoliata*, *Citrus sunki* and their hybrids under infection of *Phytophthora parasitica*," *BMC Genomics* 12(1), Article number 39. DOI: 10.1186/1471-2164-12-39
- Boland, G. J., and Hall, R. (1994). "Index of plant hosts of *Sclerotinia sclerotiorum*," *Canadian Journal of Plant Pathology* 16(2), 93-108. DOI: 10.1080/07060669409500766
- Borges, A., Ferreira, C., Saavedra, M. J., and Simoes, M. (2013). "Antibacterial activity and mode of action of ferulic and gallic acids against pathogenic bacteria," *Microbial Drug Resistance* 19(4), 256-265. DOI: 10.1089/mdr.2012.0244
- Borja, R., Martin, A., Maestro, R., Luque, M., and Durán, M. M. (1993). "Enhancement of the anaerobic digestion of wine distillery wastewater by the removal of phenolic inhibitors," *Bioresource Technology* 45(2), 99-104. DOI: 10.1016/0960-8524(93)90097-U
- Bostyn, S., Cagnon, B., and Fauduet, H. (2009). "Optimization by the simplex method of the separation of phenolic acids by high-performance liquid chromatography in wastewater olive and sugar beet vinasse," *Talanta* 80(1), 1-7. DOI: 10.1016/j.talanta.2009.06.040
- Butu, M., Stef, R., Grozea, I., Corneanu, M., and Butnariu, M. (2020). "Biopesticides: Clean and viable technology for healthy environment," in: *Bioremediation and Biotechnology*, K. Hakeem, R. Bhat, and H. Qadri (eds.), Springer, Cham, Switzerland, pp. 107-151. DOI: 10.1007/978-3-030-35691-0_6
- Cabral, J. R., Freitas, P. S. L., Bertonha, A., and Muniz, A. S. (2010). "Effects of wastewater from a cassava industry on soil chemistry and crop yield of lopsided oats (*Avena strigosa* Schreb.)," *Brazilian Archives of Biology and Technology* 53(1), 19-26. DOI: 10.1590/S1516-89132010000100003
- Cämmerer, B., and Kroh, L. W. (1995). "Investigation of the influence of reaction conditions on the elementary composition of melanoidins," *Food Chemistry* 53(1), 55-59. DOI: 10.1016/0308-8146(95)95786-6
- Cao, Y., Wang, J., Wang, Q., Liu, J., Liu, T., Knapp, C. W., and Wang, Y. (2019). "Effect of β -glycosidase supplementation on vinasse saccharification and L-lactic

- acid fermentation,” *BioResources* 14(1), 1379-1389. DOI: 10.15376/biores.14.1.1379-1389
- Capinera, J. (2020). “Introduction,” in: *Handbook of Vegetable Pests*, N. Maragioglio (ed.), Academic Press, London, United Kingdom, pp. 1-21.
- Castelian, A., de Fatima Martins, L., Cardoso, J. F. M., Silva, M. S. O., da Silva, R. S. A., Chacon-Orozco, J. G., Casteliani, A. G. B., Půža, V., Harakava, R., and Leite, L. G. (2020). “Behavioral aspects of *Sphenophorus levis* (Coleoptera: Curculionidae), damage to sugarcane and its natural infection by *Steinernema carpocapsae* (Nematoda: Rhabditidae),” *Crop Protection* 137, Article ID 105262. DOI: 10.1016/j.cropro.2020.105262
- Çelik, G., Saygın, Ö., and Balcıoğlu, I. A. (2020). “Multistage recovery process of phenolic antioxidants with a focus on hydroxytyrosol from olive mill wastewater concentrates,” *Separation and Purification Technology* 259, Article ID 117757. DOI: 10.1016/j.seppur.2020.117757
- Chand, S. K., Nanda, S., Rout, E., Mohanty, J., Mishra, R., and Joshi, R. K. (2016). “Identification and characterization of microRNAs in turmeric (*Curcuma longa* L.) responsive to infection with the pathogenic fungus *Pythium aphanidermatum*,” *Physiological and Molecular Plant Pathology* 93, 119-128. DOI: 10.1016/j.pmpp.2016.01.010
- Chen, Y., and Dai, G. (2015). “Acaricidal, repellent, and oviposition-deterrent activities of 2,4-di-tert-butylphenol and ethyl oleate against the carmine spider mite *Tetranychus cinnabarinus*,” *Journal of Pest Science* 88(3), 645-655. DOI: 10.1007/s10340-015-0646-2
- Cheng, Z., Yang, R., Wang, B., and Yang, F. (2015). “Chlorophenol degradation in papermaking wastewater through a heterogeneous ozonation process catalyzed by Fe-Mn/sepiolite,” *BioResources* 10(3), 5503-5514. DOI: 10.15376/biores.10.3.5503-5514
- Collange, B., Navarrete, M., Peyre, G., Mateille, T., and Tchamitchian, M. (2011). “Root-knot nematode (*Meloidogyne*) management in vegetable crop production: The challenge of an agronomic system analysis,” *Crop Protection* 30(10), 1251-1262. DOI: 10.1016/j.cropro.2011.04.016
- Couallier, E. M., Ruiz, B. S., Lameloise, M. L., and Decloux, M. (2006). “Usefulness of reverse osmosis in the treatment of condensates arising from the concentration of distillery vinasses,” *Desalination* 196(1-3), 306-317. DOI: 10.1016/j.desal.2006.02.002
- Dadther-Huaman, H., Machaca-Paccara, A., and Quispe-Castro, R. (2020). “Effectiveness of nine control methods of *Oregmomyza peruviana* (Granara de Willink and Diaz) (Hemiptera: Coccoidea: Eriococcidae) in *Vitis vinifera* L. ‘Negra Criolla’ and ‘Quebranta’,” *Scientia Agropecuaria* 11(1), 95-103. DOI: 10.17268/sci.agropecu.2020.01.11
- Daneshfar, A., Ghaziaskar, H. S., and Homayoun, N. (2008). “Solubility of gallic acid in methanol, ethanol, water, and ethyl acetate,” *Journal of Chemical and Engineering Data* 53(3), 776-778. DOI: 10.1021/je700633w
- Daragon, G., Trouve, G., Schönnenbeck, C., and Leyssens, G. (2014). “Recovery of an agro-industrial vinasse by adsorption on different wood materials: Parametric study at laboratory scale,” *BioResources* 9(4), 7764-7781. DOI: 10.15376/biores.9.4.7984-8001

- De León Rodríguez, A., Escalante Minakata, M. D. P., Jiménez García, M. I., Ordoñez Acevedo, L. G., Flores Flores, J. L., and Barba de la Rosa, A. P. (2008). "Characterization of volatile compounds from ethnic agave alcoholic beverages by gas chromatography-mass spectrometry," *Food Technol. Biotechnol.* 46 (4) 448–455.
- Debo, A., Yangui, T., Dhoub, A., Ksantini, M., and Sayadi, S. (2011). "Efficacy of a hydroxytyrosol-rich preparation from olive mill wastewater for control of olive psyllid, *Euphyllura olivina*, infestations," *Crop Protection* 30(12), 1529-1534. DOI: 10.1016/j.cropro.2011.08.006
- Deepak, P., Balamuralikrishnan, B., Park, S., Sowmiya, R., Balasubramani, G., Aiswarya, D., Amutha, V., and Perumal, P. (2019). "Phytochemical profiling of marine red alga, *Halymenia palmata* and its bio-control effects against Dengue Vector, *Aedes aegypti*," *South African Journal of Botany* 121, 257-266. DOI: 10.1016/j.sajb.2018.11.011
- Dekebo, A., and Jung, C. (2020). "Olfactory responses of *Aethina tumida* Murray (Coleoptera: Nitidulidae) to some major volatile compounds from hive materials and workers of *Apis mellifera*," *Journal of Asia-Pacific Entomology* 23(2), 504-508. DOI: 10.1016/j.aspen.2020.03.002
- Demuner, I. F., Colodette, J. L., Demuner, A. J., and Jardim, C. M. (2019). "Biorefinery review: Wide-reaching products through kraft lignin," *BioResources* 14(3), 7543-7581. DOI: 10.15376/biores.14.3.Demuner
- Deng, J., Xu, Z., Xiang, C., Liu, J., Zhou, L., Li, T., Yang, Z., and Ding, C. (2017). "Comparative evaluation of maceration and ultrasonic-assisted extraction of phenolic compounds from fresh olives," *Ultrasonics Sonochemistry* 37, 328-334. DOI: 10.1016/j.ultsonch.2017.01.023
- Deng, N., Li, M., Zhao, L., Lu, C., de Rooy, S. L., and Warner, I. M. (2011). "Highly efficient extraction of phenolic compounds by use of magnetic room temperature ionic liquids for environmental remediation," *Journal of Hazardous Materials* 192(3), 1350-1357. DOI: 10.1016/j.jhazmat.2011.06.053
- Dharni, S., Sengupta, S., Maurya, A., Samad, A., Srivastava, S. K., Sharma, A., and Patra, D. D. (2014). "Purification, characterization, and *in vitro* activity of 2,4-di-tert-butylphenol from *Pseudomonas monteilii* PsF84: Conformational and molecular docking studies," *Journal of Agricultural and Food Chemistry* 62(26), 6138-6146. DOI: 10.1021/jf5001138
- Di Ilio, V., and Cristofaro, M. (2020). "Polyphenolic extracts from the olive mill wastewater as a source of biopesticides and their effects on the life cycle of the Mediterranean fruit fly *Ceratitis capitata* (Diptera, Tephritidae)," *International Journal of Tropical Insect Science Online*, 1-8. DOI: 10.1007/s42690-020-00224-6
- Di Mauro, M. D., Giardina, R. C., Fava, G., Mirabella, E. F., Acquaviva, R., Renis, M., and D'antona, N. (2017). "Polyphenolic profile and antioxidant activity of olive mill wastewater from two Sicilian olive cultivars: Cerasuola and Nocellara etnea," *European Food Research and Technology* 243(11), 1895-1903. DOI: 10.1007/s00217-017-2893-3
- Díaz, B., Gomes, A., Freitas, M., Fernandes, E., Nogueira, D. R., González, J., Moure, A., Levoso, A., Vinardell, M. P., Mitjans, M., Domínguez, H., and Parajó, J. C. (2012). "Valuable polyphenolic antioxidants from wine vinasses," *Food and Bioprocess Technology* 5(7), 2708-2716. DOI: 10.1007/s11947-011-0569-8
- do Carmo, J. B., Filoso, S., Zotelli, L. C., de Sousa Neto, E. R., Pitombo, L. M., Duarte-Neto, P. J., Vargas, V. P., Andrade, C. A., Gava, G. J. C., Rossetto, R., *et al.* (2012).

- “Infield greenhouse gas emissions from sugarcane soils in Brazil: Effects from synthetic and organic fertilizer application and crop trash accumulation,” *GCB Bioenergy* 5(3), 267-280. DOI: 10.1111/j.1757-1707.2012.01199.x
- dos Santos Moura, A., de Medeiros, E. V., da Silva Oliveira, J. E., da Franca, R. F., Lira, A. D., Martins Filho, A. P., and Badji, C. A. (2018). “Does cassava wastewater with a short incubation time affect soil organic carbon, microbial community and enzymatic activities?,” *CATENA* 163, 354-360. DOI: 10.1016/j.catena.2018.01.001
- EFSA Panel on Plant Health PLH (2014). “Scientific opinion on the pest categorisation of *Verticillium dahliae* Kleb,” *EFSA Journal* 12(12), article no. 3928. DOI: 10.2903/j.efsa.2014.3928
- El Hassani, F. Z., Fadile, A., Faouzi, M., Zinedine, A., Merzouki, M., and Benlemlih, M. (2020). “The long term effect of olive mill wastewater (OMW) on organic matter humification in a semi-arid soil,” *Heliyon* 6(1), article ID e03181. DOI: 10.1016/j.heliyon.2020.e03181
- El-Abbassi, A., Saadaoui, N., Kiai, H., Raiti, J., and Hafidi, A. (2017). “Potential applications of olive mill wastewater as biopesticide for crops protection,” *Science of The Total Environment* 576, 10-21. DOI: 10.1016/j.scitotenv.2016.10.032
- El-Sayed, A. S. A., and Ali, G. S. (2020). “*Aspergillus flavipes* is a novel efficient biocontrol agent of *Phytophthora parasitica*,” *Biological Control* 140, article ID 104072. DOI: 10.1016/j.biocontrol.2019.104072
- Erb, M., Meldau, S., and Howe, G. A. (2012). “Role of phytohormones in insect-specific plant reactions,” *Trends in Plant Science* 17(5), 250-259. DOI: 10.1016/j.tplants.2012.01.003
- Erukainure, O. L., Sanni, O., and Islam, M. S. (2018). “*Clerodendrum volubile*: Phenolics and applications to health,” in: *Polyphenols: Mechanisms of Action in Human Health and Disease*, R. R. Watson, V. R. Preedy, and S. Zibadi (eds.), Academic Press, Cambridge, MA, USA, pp. 53-68. DOI: 10.1016/B978-0-12-813006-3.00006-4
- Espinoza-Acosta, J. L., Torres-Chávez, P. I., Ramírez-Wong, B., López-Saiz, C. M., and Montaña-Leyva, B. (2016). “Antioxidant, antimicrobial, and antimutagenic properties of technical lignins and their applications,” *BioResources* 11(2), 5452-5481. DOI: 10.15376/biores.11.2.Espinoza_Acosta
- Félix, E., Contreras-Ramos, S. M., Davila-Vazquez, G., Rodríguez-Campos, J., and Marino, E. (2018). “Identification and quantification of volatile compounds found in vinasses from two different processes of tequila production,” *Energies* 11(3), Article Number 490. DOI: 10.3390/en11030490
- Fernandes, J. M. C., Sousa, R. M. O. F., Fraga, I., Sampaio, A., Amaral, C., Bezerra, R. M. F., and Dias, A. A. (2020). “Fungal biodegradation and multi-level toxicity assessment of vinasse from distillation of winemaking by-products,” *Chemosphere* 238, Article ID 124572. DOI: 10.1016/j.chemosphere.2019.124572
- Freitas, P. V., da Silva, D. R., Beluomini, M. A., da Silva, J. L., and Stradiotto, N. R. (2018). “Determination of phenolic acids in sugarcane vinasse by HPLC with pulse amperometry,” *Journal of Analytical Methods in Chemistry* 2018, Article ID 4869487. DOI: 10.1155/2018/4869487
- Fuess, L., Garcia, M. L., and Zaiat, M. (2018). “Seasonal characterization of sugarcane vinasse: Assessing environmental impacts from fertirrigation and the bioenergy recovery potential through biodigestion,” *Science of The Total Environment* 634, Article ID 4686011. DOI: 10.1016/j.scitotenv.2018.03.326

- Gonzalez-Coloma, A., Reina, M., Díaz, C., Fraga, B. M., and Santana-Meridas, O. (2013). "Natural product-based biopesticides for insect control" in: *Chemistry, Molecular Sciences and Chemical Engineering*, Elsevier, Amsterdam, Netherlands, pp. 237-268. DOI: 10.1016/B978-0-12-409547-2.02770-0
- Gu, S., Choi, J. W., Lee, H., Suh, D. J., Choi, J., and Ha, J. M. (2020). "Improved catalytic depolymerization of lignin waste using carbohydrate derivatives," *Environmental Pollution* 268(Part A), article ID 115674. DOI: 10.1016/j.envpol.2020.115674
- Hahn, M., Viaud, M., and van Kan, J. (2014). "The genome of *Botrytis cinerea*, a ubiquitous broad host range necrotroph," in: *Genomics of Plant-associated Fungi and Oomycetes: Dicot Pathogens*, Springer, Berlin, Germany pp. 19-44. DOI: 10.1007/978-3-662-44056-8_2
- Hou, Y. T., Ren, J., and Liu, H. J. (2012). "Efficiency of electrolyzed water (EW) on inhibition of *Phytophthora parasitica* var. *nicotianae* growth *in vitro*," *Crop Protection* 42, 128-133. DOI: 10.1016/j.cropro.2012.06.014
- Hougardy, E., Wang, X., Hogg, B. N., Johnson, M. W., Daane, K. M., and Pickett, C. H. (2020). "Current distribution of the olive psyllid, *Euphyllura olivina*, in California and initial evaluation of the Mediterranean parasitoid *Psyllaephagus euphyllurae* as a biological control candidate," *Insects* 11(3), article no. 146. DOI: 10.3390/insects11030146
- Huang, X., Zhao, J., Zhou, X., Zhang, J., and Cai, Z. (2019). "Differential responses of soil bacterial community and functional diversity to reductive soil disinfection and chemical soil disinfection," *Geoderma* 348, 124-134. DOI: 10.1016/j.geoderma.2019.04.027
- Jactel, H., Koricheva, J., and Castagnyrol, B. (2019). "Responses of forest insect pests to climate change: Not so simple," *Current Opinion in Insect Science* 35, 103-108. DOI: 10.1016/j.cois.2019.07.010
- Jelvez Serra, N. S., Goulart, H. F., Triana, M. F., Dos Santos Tavares, S., Almeida, C. I. M., Da Costa, J. G., Santana, A. E. G., and Zhu, J. J. (2017). "Identification of stable fly attractant compounds in vinasse, a byproduct of sugarcane-ethanol distillation," *Medical and Veterinary Entomology* 31(4), 381-391. DOI: 10.1111/mve.12246
- Jinxiang, C., Yang, J., Lanlan, M., Li, J., Nasir, S., and Kyung, K. C. (2020). "Structure-antioxidant activity relationship of methoxy, phenolic hydroxyl, and carboxylic acid groups of phenolic acids," *Scientific Reports (Nature Publisher Group)* 10(1), article no. 2611. DOI: 10.1038/s41598-020-59451-z
- Jung, K. H., You, S. K., Moon, S. K., and Lee, U. S. (2007). "Furfural from pine needle extract inhibits the growth of a plant pathogenic fungus, *Alternaria mali*," *Mycobiology* 35(1), 39-43. DOI: 10.4489/myco.2007.35.1.039
- Kannabiran, K. (2016). "Bioactivity of pyrrolo [1, 2-a] pyrazine-1, 4-dione, hexahydro-3-(phenylmethyl)-extracted from *Streptomyces* sp. VITPK9 isolated from the salt spring habitat of Manipur, India," *Asian Journal of Pharmaceutics* 10(4), 265-270. DOI: 10.22377/ajp.v10i04.865
- Kaushik, A., Basu, S., Batra, V. S., and Balakrishnan, M. (2018). "Fractionation of sugarcane molasses distillery wastewater and evaluation of antioxidant and antimicrobial characteristics," *Industrial Crops and Products* 118, 73-80. DOI: 10.1016/j.indcrop.2018.03.040

- Khan, M. F., and Murphy, C. D. (2020). “3-Hydroxytyrosol regulates biofilm growth in *Cunninghamella elegans*,” *Fungal Biology In Press*, Corrected proof available online. DOI: 10.1016/j.funbio.2020.10.011
- Kim, D. (2018). “Physico-chemical conversion of lignocellulose: Inhibitor effects and detoxification strategies: A mini review,” *Molecules* 23(2), Article number 309. DOI: 10.3390/molecules23020309
- Kim, J. H., Chan, K. L., Mahoney, N., and Campbell, B. C. (2011). “Antifungal activity of redox-active benzaldehydes that target cellular antioxidation,” *Annals of Clinical Microbiology and Antimicrobials* 10(1), article no. 23. DOI:10.1186/1476-0711-10-23
- Kotowska, U., Kapelewska, J., and Sturgulewska, J. (2014). “Determination of phenols and pharmaceuticals in municipal wastewaters from Polish treatment plants by ultrasound-assisted emulsification–microextraction followed by GC–MS,” *Environmental Science and Pollution Research* 21(1), 660-673. DOI: 10.1007/s11356-013-1904-6
- Kou, H., Sun, Y., Dong, Z., and Zhang, Z. (2020). “Comparison between sustained effects of spray and injection thiamethoxam on apple aphids and non-target insects in apple orchard,” *Ecotoxicology and Environmental Safety* 207, article ID 111307. DOI: 10.1016/j.ecoenv.2020.111307
- Koul, O., and Walia, S. (2009). “Comparing impacts of plant extracts and pure allelochemicals and implications for pest control,” *CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources* 4(49), 1-30. DOI: 10.1079/PAVSNNR20094049
- Larif, M., Ouhssine, M., Soulaymani, A., and Elmidaoui, A. (2015). “Potential effluent oil mills and antibacterial activity polyphenols against some pathogenic strains,” *Research on Chemical Intermediates* 41(2), 1213-1225. DOI: 10.1007/s11164-013-1267-0
- Larif, M., Zarrouk, A., Soulaymani, A., and Elmidaoui, A. (2013). “New innovation in order to recover the polyphenols of olive mill wastewater extracts for use as a biopesticide against the *Euphyllura olivina* and *Aphis citricola*,” *Research on Chemical Intermediates* 39(9), 4303-4313. DOI: 10.1007/s11164-012-0947-5
- Lebeda, A., Mieslerová, B., Jankovics, T., Kiss, L., and Van der Linde, E. J. (2015). “First detection of tomato powdery mildew caused by *Oidium neolycopersici* in South Africa,” *South African Journal of Botany* 99, 153-157. DOI: 10.1016/j.sajb.2015.03.196
- Lengai, G. M. W., Muthomi, J. W., and Mbega, E. R. (2020). “Phytochemical activity and role of botanical pesticides in pest management for sustainable agricultural crop production,” *Scientific African* 7, Article ID e00239. DOI: 10.1016/j.sciaf.2019.e00239
- Lin, S., Tang, T., Cang, T., Yu, S., Ying, Z., Gu, S., and Zhang, Q. (2020). “The distributions of three fungicides in vegetables and their potential health risks in Zhejiang, China: A 3-year study (2015–2017),” *Environmental Pollution* 267, 115481. DOI: 10.1016/j.envpol.2020.115481
- Lind, H., Jonsson, H., and Schnürer, J. (2005). “Antifungal effect of dairy propionibacteria—contribution of organic acids,” *International Journal of Food Microbiology* 98(2), 157-165. DOI: 10.1016/j.ijfoodmicro.2004.05.020
- Liu, J., Wang, Q., Wang, S., Sun, X., Ma, H., and Tushiro, Y. (2013). “Effects of pretreatment on the microbial community and l-lactic acid production in vinasse

- fermentation,” *Journal of biotechnology* 164(2), 260-265. DOI: 10.1016/j.jbiotec.2012.08.014
- Logan, J. A., Régnière, J., and Powell, J. A. (2003). “Assessing the impacts of global warming on forest pest dynamics,” *Frontiers in Ecology and the Environment* 1(3), 130-137. DOI: 10.1890/1540-9295(2003)001[0130:ATIOWG]2.0.CO;2
- Mansfield, J., Genin, S., Magori, S., Citovsky, V., Sriariyanum, M., Ronald, P., Dow, M., Verdier, V., Beer, S. V., Machado, M. A., *et al.* (2012). “Top 10 plant pathogenic bacteria in molecular plant pathology,” *Molecular Plant Pathology* 13(6), 614-629. DOI: 10.1111/j.1364-3703.2012.00804.x
- Markakis, E. A., Fountoulakis, M. S., Daskalakis, G. C., Kokkinis, M., and Ligoxigakis, E. K. (2016). “The suppressive effect of compost amendments on *Fusarium oxysporum* f. sp. *radicis-cucumerinum* in cucumber and *Verticillium dahliae* in eggplant,” *Crop Protection* 79, 70-79. DOI: 10.1016/j.cropro.2015.10.015
- Martínez, L., Ros, G., and Nieto, G. (2018). “Hydroxytyrosol: Health benefits and use as functional ingredient in meat,” *Medicines* 5(1), article no. 13. DOI: 10.3390/medicines5010013
- Martins, L. F., Tonelli, M., Bento, J. M. S., Bueno, C. J., and Leite, L. G. (2020). “Attraction of the sugarcane billbug, *Sphenophorus levis*, to vinasse and its volatile composition,” *Chemoecology* 30, 205-214. DOI: 10.1007/s00049-020-00310-8
- Matthews, G. A. (2018). “Pesticides in the early part of the 20th century,” in: *A History of Pesticides*, CABI, London, United Kingdom, pp. 1-27.
- Meissle, M., Mouron, P., Musa, T., Bigler, F., Pons, X., Vasileiadis, V. P., Otto, S., Antichi, D., Kiss, J., Pálincás, J., *et al.* (2010). “Pests, pesticide use and alternative options in European maize production: Current status and future prospects,” *Journal of Applied Entomology* 134(5), 357-375. DOI: 10.1111/j.1439-0418.2009.01491.x
- Mekki, A., Dhouib, A., and Sayadi, S. (2013). “Review: Effects of olive mill wastewater application on soil properties and plants growth,” *International Journal of Recycling of Organic Waste in Agriculture* 2(1), 1-7. DOI: 10.1186/2251-7715-2-15
- Meng, Y., Zhang, Q., Ding, W., and Shan, W. (2014). “*Phytophthora parasitica*: A model oomycete plant pathogen,” *Mycology* 5(2), 43-51. DOI: 10.1080/21501203.2014.917734
- Minaeimoghadam, M., Askarianzadeh, A., Imani, S., Shojaei, M., Larijani, K., and Abbasipour, H. (2017). “Identification of chemical compounds of the pheromone in different ages of female adults of the clearwing moth, *Paranthrene diaphana* Dalla Torre and Strand,” *Archives of Phytopathology and Plant Protection* 50(19-20), 1019-1033. DOI: 10.1080/03235408.2017.1411174
- Mohamed, H. A., Nasr, N. F., and Abdelkreem, K. I. (2015). “Management of tomato damping-off disease caused by *Fusarium oxysporum* and *Rhizoctonia solani* using chemical and biological degradable olive mill waste water,” *Egyptian Journal of Biological Pest Control* 25(2), Article number 367-377.
- Molinari, S. (2011). “Natural genetic and induced plant resistance, as a control strategy to plant-parasitic nematodes alternative to pesticides,” *Plant Cell Reports* 30(3), 311-323. DOI: 10.1007/s00299-010-0972-z
- Neves, O. S. C., Souza, A. S., Costa, M. A., Sousa, L. D. A., Viana, A. E. S., and Neves, V. B. F. (2014). “Persistência do cianeto e estabilização do pH em manipueira,” *Revista Brasileira de Tecnologia Agroindustrial* 8(1), 1274-1284. DOI: 10.3895/S1981-36862014000100012

- Noushini, S., Perez, J., Park, S. J., Holgate, D., Jamie, I., Jamie, J., and Taylor, P. (2020). "Rectal gland chemistry, volatile emissions, and antennal responses of male and female banana fruit fly, *Bactrocera musae*," *Insects* 11(1), article no. 32. DOI: 10.3390/insects11010032
- Ntalli, N., Adamski, Z., Doula, M., and Monokrousos, N. (2020). "Nematicidal amendments and soil remediation," *Plants* 9(4), 429. DOI: 10.3390/plants9040429
- Núñez-Zofío, M., Larregla, S., Garbisu, C., Guerrero, M. M., Lacasa, C. M., and Lacasa, A. (2013). "Application of sugar beet vinasse followed by solarization reduces the incidence of *Meloidogyne incognita* in pepper crops while improving soil quality," *Phytoparasitica* 41(2), 181-191. DOI: 10.1007/s12600-012-0277-6
- Ordaz-Diaz, L. A., Madrid-del Palacio, M., Rodriguez-Flores, F. J., and Bailón-Salas, A. M. (2019). "Aprovechamiento de los subproductos valiosos de la industria del mezcal en Durango [Taking advantage of the valuable by-products of the mezcal industry in Durango]," in: *Colaboraciones de Cuerpos Académicos en Innovación Tecnológica [Collaborations of Academic Bodies in Technological Innovation]*, Ibero-American Network of Research Academies A.C., Veracruz, México, pp. 14-23.
- Owuor, P. O. (2003). "TEA analysis and tasting," in: *Encyclopedia of Food Sciences and Nutrition*, B. Caballero (ed.), Academic Press, Cambridge, MA, USA, pp. 1-10.
- Padmaja, G. (1995). "Cyanide detoxification in cassava for food and feed uses," *Critical Reviews in Food Science and Nutrition* 35(4), 299-339. DOI: 10.1080/10408399509527703
- Papaianni, M., Paris, D., Woo, S. L., Fulgione, A., Rigano, M. M., Parrilli, E., Tutino, M. L., Marra, R., Manganiello, G., Casillo, A., et al. (2020). "Plant dynamic metabolic response to bacteriophage treatment after *Xanthomonas campestris* pv. *campestris* infection," *Frontiers in Microbiology* 11, Article Number 732. DOI: 10.3389/fmicb.2020.00732
- Paslaru, A. I., Verhulst, N. O., Maurer, L. M., Brendle, A., Pauli, N., Vöglin, A., Renzullo, S., Ruedin, Y., Hoffmann, B., Torgerson, P. R., et al. (2020). "Potential mechanical transmission of Lumpy skin disease virus (LSDV) by the stable fly (*Stomoxys calcitrans*) through regurgitation and defecation," *Current Research in Insect Science* 1, Article ID 100007. DOI: 10.1016/j.cris.2020.100007
- Patzke, H., and Schieber, A. (2018). "Growth-inhibitory activity of phenolic compounds applied in an emulsifiable concentrate-ferulic acid as a natural pesticide against *Botrytis cinerea*," *Food Research International* 113, 18-23. DOI: 10.1016/j.foodres.2018.06.062
- Paulitz, T. C., and Schroeder, K. L. (2005). "A new method for the quantification of *Rhizoctonia solani* and *R. oryzae* from soil," *Plant Disease* 89(7), 767-772. DOI: 10.1094/PD-89-0767
- Peng, C., Zhang, G., Han, J., and Li, X. (2019). "Hydrothermal conversion of lignin and black liquor for phenolics with the aids of alkali and hydrogen donor," *Carbon Resources Conversion* 2(2), 141-150. DOI: 10.1016/j.crcon.2019.06.004
- Pereira, C. S. M., Silva, V. M. T. M., and Rodrigues, A. E. (2011). "Ethyl lactate as a solvent: Properties, applications and production processes—A review," *Green Chemistry* 13(10), 2658-2671. DOI: 10.1039/c1gc15523g
- Phanapavudhikul, S. (1999). "Direct use of spent distillery wash liquor on paddy fields in Thailand," *Water and Environment Journal* 13(6), 420-422. DOI: 10.1111/j.1747-6593.1999.tb01079.x

- Phillips, C. L., and Nickerson, N. (2015). "Soil respiration," in: *Reference Module in Earth Systems and Environmental Sciences*, S. Elias (ed.), Elsevier, Amsterdam, Netherlands, pp. 515-527. DOI: 10.1016/B978-0-12-409548-9.09442-2
- Piśyć, S., Natorff, R., Gawińska-Urbanowicz, H., and Kruszewska, J. S. (2015). "Fusarium sambucinum astA gene expressed during potato infection is a functional orthologue of Aspergillus nidulans astA," *Fungal Biology* 119(6), 509-517. DOI: 10.1016/j.funbio.2015.02.002
- Pinto-Zevallos, D. M., Pereira-Querol, M., and Ambrogi, B. G. (2018). "Cassava wastewater as a natural pesticide: Current knowledge and challenges for broader utilization," *Annals of Applied Biology* 173(3), 191-201. DOI: 10.1111/aab.12464
- Ponte, J. D., Franco, A., and Santos, J. H. R. D. (1988). "Teste preliminar sobre a utilização da manipueira como inseticida [Preliminary test on the use of manipueira as an insecticide]," *Revista Brasileira de Mandioca* 7, 89-90.
- Ponte, J., and Franco, A. (1983). "Influencia da idade da manipueira na preservacao do potencial nematicida do composto (nota previa) [Influence of the age of the manipueira on the preservation of the nematicidal potential of the compound (previous note)]," *Brazilian Society of Nematology* 7, article ID 35215.
- Puoci, F., Scoma, A., Cirillo, G., Bertin, L., Fava, F., and Picci, N. (2012). "Selective extraction and purification of gallic acid from actual site olive mill wastewaters by means of molecularly imprinted microparticles," *Chemical Engineering Journal* 198-199, 529-535. DOI: 10.1016/j.cej.2012.05.095
- Rashwan, R. S., and Hammad, D. M. (2020). "Toxic effect of Spirulina platensis and Sargassum vulgare as natural pesticides on survival and biological characteristics of cotton leaf worm Spodoptera littoralis," *Scientific African* 8, Article ID e00323. DOI: 10.1016/j.sciaf.2020.e00323
- Ravindran, D. R., Bharathithasan, M., Ramaiah, P., Rasat, M. S. M., Rajendran, D., Srikumar, S., Ishak, I. H., Said, A. R., Ravi, R., and Amin, M. F. M. (2020). "Chemical composition and larvicidal activity against Aedes mosquitos of flower extracts from Clitoria ternatea," *Journal of Chemistry* 2020, 1-9. DOI: 10.1101/2020.03.31.018465
- Rebollo-Romero, I., Fernández-Cruz, E., Carrasco-Galán, F., Valero, E., Cantos-Villar, E., Cerezo, A. B., Troncoso, A. M., and Garcia-Parrilla, M. C. (2020). "Factors influencing the production of the antioxidant hydroxytyrosol during alcoholic fermentation: Yeast strain, initial tyrosine concentration and initial must," *LWT* 130, Article ID 109631. DOI: 10.1016/j.lwt.2020.109631
- Reddy, N., and Yang, Y. (2005). "Biofibers from agricultural byproducts for industrial applications," *Trends in Biotechnology* 23(1), 22-27. DOI: 10.1016/j.tibtech.2004.11.002
- Regmi, H., and Desaegeer, J. (2020). "Integrated management of root-knot nematode (Meloidogyne spp.) in Florida tomatoes combining host resistance and nematicides," *Crop Protection* 134, article ID 105170. DOI: 10.1016/j.cropro.2020.105170
- Robles-González, V., Galíndez-Mayer, J., Rinderknecht-Seijas, N., and Poggi-Varaldo, H. M. (2012). "Treatment of mezcal vinasses: A review," *Journal of Biotechnology* 157(4), 524-546. DOI: 10.1016/j.jbiotec.2011.09.006
- Rosazza, J. P. N., Huang, Z., Dostal, L., Volm, T., and Rousseau, B. (1995). "Review: Biocatalytic transformations of ferulic acid: An abundant aromatic natural product," *Journal of Industrial Microbiology* 15(6), 457-471. DOI: 10.1007/BF01570016

- Roy, R., Jadhav, B., Rahman, S., and Raynie, D. E. (2020). "Characterization of residue from catalytic hydrothermal depolymerization of lignin," *Current Research in Green and Sustainable Chemistry* 4, article ID 100052. DOI: 10.1016/j.crgsc.2020.100052
- Rubin-Aguirre, A., Sáenz-Romero, C., Lindig-Cisneros, R., del-Rio-Mora, A. A., Tena-Morelos, C. A., Campos-Bolaños, R., and del-Val, E. (2015). "Bark beetle pests in an altitudinal gradient of a Mexican managed forest," *Forest Ecology and Management* 343, 73-79. DOI: 10.1016/j.foreco.2015.01.028
- Rubio-Senent, F., Fernández-Bolaños, J., García-Borrego, A., Lama-Muñoz, A., and Rodríguez-Gutiérrez, G. (2017). "Influence of pH on the antioxidant phenols solubilised from hydrothermally treated olive oil by-product (alperujo)," *Food Chemistry* 219, 339-345. DOI: 10.1016/j.foodchem.2016.09.141
- Sadaria, A. M., Labban, C. W., Steele, J. C., Maurer, M. M., and Halden, R. U. (2019). "Retrospective nationwide occurrence of fipronil and its degradates in US wastewater and sewage sludge from 2001-2016," *Water Research* 155, 465-473. DOI: 10.1016/j.watres.2019.02.045
- Saeed, Q., Ahmad, F., Iqbal, N., and Zaka, S. M. (2019). "Chemical control of polyphagous pests on their auxiliary hosts can minimize insecticide resistance: A case study of *Spodoptera exigua* Hübner (Lepidoptera: Noctuidae) in cotton agroecosystem," *Ecotoxicology and Environmental Safety* 171, 721-727. DOI: 10.1016/j.ecoenv.2019.01.038
- Sanchez, S., Valadez-Rosales, M. F., Correa-Camacho, J. M., and Marino, E. (2019). "Recovery of phenolic compounds from Tequila vinasses using adsorption resins in two configurations (batch and continuous)," *Journal of Chemical and Physical Energy* 6(20), 1-6. DOI: 10.35429/JCPE.2019.20.6.1.6
- Santos, A., and Ponte, J. D. (1993). *Ação Fungicida da Manipueira No Controle de Oídio [Fungicidal Action of Manipueira in Powdery Mildew Control]*, Fitopatol, Lavras, Brazil, pp. 18, 302.
- Santos, M., Diánez, F., De Cara, M., and Tello, J. C. (2008). "Possibilities of the use of vinasses in the control of fungi phytopathogens," *Bioresource Technology* 99(18), 9040-9043. DOI: 10.1016/j.biortech.2008.04.032
- Sarip, H., Hossain, S., Azemi, M., and Allaf, K. (2016). "A review of the thermal pretreatment of lignocellulosic biomass towards glucose production: Autohydrolysis with DIC technology," *BioResources* 11(4), 10625-10653. DOI: 10.15376/biores.11.4.Sarip
- Sassi, A. B., Boularbah, A., Jaouad, A., Walker, G., and Boussaid, A. (2006). "A comparison of olive oil mill wastewaters (OMW) from three different processes in Morocco," *Process Biochemistry* 41(1), 74-78. DOI: 10.1016/j.procbio.2005.03.074
- Savary, S., Willocquet, L., Pethybridge, S. J., Esker, P., McRoberts, N., and Nelson, A. (2019). "The global burden of pathogens and pests on major food crops," *Nature Ecology and Evolution* 3, 430-439. DOI: 10.1038/s41559-018-0793-y
- Saxena, S., and Strobel, G. A. (2020). "*Marvellous Muscodor* spp.: Update on their biology and applications," *Microbial Ecology* 24, 1-16. DOI: 10.1007/s00248-020-01644-0
- Sedenho, G. C., Da Silva, J. L., Beluomini, M. A., De Sá, A. C., and Stradiotto, N. R. (2017). "Determination of electroactive organic acids in sugarcane vinasse by high performance anion-exchange chromatography with pulsed amperometric detection using a nickel nanoparticle modified boron-doped diamond," *Energy & Fuels* 31(3), 2865-2870. DOI: 10.1021/acs.energyfuels.6b02783

- Shang, X., Dai, L., Yang, C., Guo, X., Liu, Y., Miao, X., and Zhang, J. (2020). "A value-added application of eugenol as acaricidal agent: The mechanism of action and the safety evaluation," *Journal of Advanced Research*, In Press, Corrected proof available online. DOI: 10.1016/j.jare.2020.12.010
- Sharma, A., Shukla, A., Attri, K., Kumar, M., Kumar, P., Suttee, A., Singh, G., Pratap-Barnwal, R. P., and Singla, N. (2020). "Global trends in pesticides: A looming threat and viable alternatives," *Ecotoxicology and Environmental Safety* 201, Article ID 110812. DOI: 10.1016/j.ecoenv.2020.110812
- Simon-Delso, N., Amaral-Rogers, V., Belzunces, L. P., Bonmatin, J. M., Chagnon, M., Downs, C., Furlan, L., Gibbons, D. W., Giorio, C., Girolami, V., *et al.* (2014). "Systemic insecticides (neonicotinoids and fipronil): Trends, uses, mode of action and metabolites," *Environmental Science and Pollution Research* 22(1), 5-34. DOI: 10.1007/s11356-014-3470-y
- Sleven, R., Lanckacker, E. A., Delputte, P. L., Maes, L., and Cos, P. (2016). "Evaluation of topical antifungal products in an *in vitro* onychomycosis model," *Mycoses* 59(5), 327-330. DOI: 10.1111/myc.12475
- Soobadar, A., and Ng Kee Kwong, K. R. (2012). "Impact of high rates of vinasse on some pertinent soil characteristics and on sugarcane yield in Mauritius," *Journal of Sustainable Agriculture* 36(1), 36-53. DOI: 10.1080/10440046.2011.620226
- Swett, C. L., Butler, B. B., Peres, N. A., Koivunen, E. E., Hellman, E. M., and Beaulieu, J. R. (2020). "Using model-based fungicide programming to effectively control Botrytis and Anthracnose fruit rots in Mid-Atlantic strawberry fields and co-manage strawberry sap beetle (*Stelidota geminata*)," *Crop Protection* 134, article no. 105175. DOI: 10.1016/j.cropro.2020.105175
- Tejada, M., García-Martínez, A. M., and Parrado, J. (2009). "Effects of a vermicompost composted with beet vinasse on soil properties, soil losses and soil restoration," *CATENA* 77(3), 238-247. DOI: 10.1016/j.catena.2009.01.004
- Thakur, N., Kaur, S., Tomar, P., Thakur, S., and Yadav, A. N. (2020). "Microbial biopesticides: Current status and advancement for sustainable agriculture and environment," in: *New and Future Developments in Microbial Biotechnology and Bioengineering*, Elsevier, Amsterdam, Netherlands, pp. 243-282. DOI: 10.1016/B978-0-12-820526-6.00016-6
- Tiwari, R. K., Kumar, R., Sharma, S., Sagar, V., Aggarwal, R., Naga, K. C., Lal, M. K., Chourasia, K. N., Kumar, D., and Kumar, M. (2020). "Potato dry rot disease: Current status, pathogenomics and management," *3 Biotech* 10(11), Article Number 503. DOI: 10.1007/s13205-020-02496-8
- Uğurlu, M., and Kula, I. (2007). "Decolourization and removal of some organic compounds from olive mill wastewater by advanced oxidation processes and lime treatment," *Environmental Science and Pollution Research-International* 14(5), 319-325. DOI: 10.1065/espr2006.06.315
- Ullah, I., Khan, A. L., Ali, L., Khan, A. R., Waqas, M., Hussain, J., Lee, I., and Shin, J. H. (2015). "Benzaldehyde as an insecticidal, antimicrobial, and antioxidant compound produced by *Photobacterium temperata* M1021," *Journal of Microbiology* 53(2), 127-133. DOI: 10.1007/s12275-015-4632-4
- Varsha, K. K., Devendra, L., Shilpa, G., Priya, S., Pandey, A., and Nampoothiri, K. M. (2015). "2,4-Di-tert-butyl phenol as the antifungal, antioxidant bioactive purified from a newly isolated *Lactococcus* sp.," *International Journal of Food Microbiology* 211, 44-50. DOI: 10.1016/j.ijfoodmicro.2015.06.025

- Vigo, C., Norman, J. R., and Hooker, J. E. (2000). "Biocontrol of the pathogen *Phytophthora parasitica* by arbuscular mycorrhizal fungi is a consequence of effects on infection loci," *Plant Pathology* 49(4), 509-514. DOI: 10.1046/j.1365-3059.2000.00473.x
- Villaverde, J. J., Sandín-España, P., Sevilla-Morán, B., López-Goti, C., and Alonso-Prados, J. L. (2016). "Biopesticides from natural products: Current development, legislative framework, and future trends," *BioResources* 11(2), 5618-5640. DOI: 10.15376/biores.11.2.Villaverde
- Villegas, L. G. C., Mashhadi, N., Chen, M., Mukherjee, D., Taylor, K. E., and Biswas, N. (2016). "A short review of techniques for phenol removal from wastewater," *Current Pollution Reports* 2(3), 157-167. DOI: 10.1007/s40726-016-0035-3
- Wang, F. (2020). "Determination of lignin concentration in bleaching effluent via Coomassie brilliant blue G-250 method," *BioResources* 15(1), 228-239. DOI: 10.15376/biores.15.1.228-239
- Wang, Y., Liu, H., Zhang, D., Liu, J., Wang, J., Wang, S., and Sun, B. (2019). "Baijiu vinasse extract scavenges glyoxal and inhibits the formation of N ϵ -carboxymethyl-lysine in dairy food," *Molecules* 24(8), Article ID 1526. DOI: 10.3390/molecules24081526
- Wathier, E., Andreani, C. L., Torres, D. G. B., Kuczman, O., Tavares, M. H. F., Lopes, D. D., and Gomes, S. D. (2019). "Cassava wastewater treatment in fixed-bed reactors: Organic matter removal and biogas production," *Frontiers in Sustainable Food Systems* 3, Article number 6. DOI: 10.3389/fsufs.2019.00006
- Wedge, R. M., Abt, E. N., and Hobbs, C. H. (2001). "Methyl bromide risk characterization in California," *Journal of Toxicology and Environmental Health Part B: Critical Reviews* 4(3), 333-339. DOI: 10.1080/10937400119036
- Wingfield, M. J., Brockerhoff, E. G., Wingfield, B. D., and Slippers, B. (2015). "Planted forest health: The need for a global strategy," *Science* 349(6250), 832-836. DOI: 10.1126/science.aac6674
- Xin, X., Kvitko, B., and He, S. Y. (2018). "*Pseudomonas syringae*: What it takes to be a pathogen," *Nature Reviews Microbiology* 16(5), 316-328. DOI: 10.1038/nrmicro.2018.17
- Yang, Z., Yu, Z., Lei, L., Xia, Z., Shao, L., Zhang, K., and Li, G. (2012). "Nematicidal effect of volatiles produced by *Trichoderma* sp.," *Journal of Asia-Pacific Entomology* 15(4), 647-650. DOI: 10.1016/j.aspen.2012.08.002
- Yangui, T., Dhouib, A., Rhouma, A., and Sayadi, S. (2009). "Potential of hydroxytyrosol-rich composition from olive mill wastewater as a natural disinfectant and its effect on seeds vigour response," *Food Chemistry* 117(1), 1-8. DOI: 10.1016/j.foodchem.2009.03.069
- Yangui, T., Sayadi, S., Rhouma, A., and Dhouib, A. (2010). "Potential use of hydroxytyrosol-rich extract from olive mill wastewater as a biological fungicide against *Botrytis cinerea* in tomato," *Journal of Pest Science* 83(4), 437-445. DOI: 10.1007/s10340-010-0314-5
- Zema, D. A., Lucas-Borja, M. E., Andiloro, S., Tamburino, V., and Zimbone, S. M. (2019). "Short-term effects of olive mill wastewater application on the hydrological and physico-chemical properties of a loamy soil," *Agricultural Water Management* 221, 312-321. DOI: 10.1016/j.agwat.2019.04.011

Zhang, M., Wang, Z., Li, P., Zhang, H., and Xie, L. (2017). "Bio-refractory dissolved organic matter and colorants in cassava distillery wastewater: Characterization, coagulation treatment and mechanisms," *Chemosphere* 178, 259-267. DOI: 10.1016/j.chemosphere.2017.03.065

Article submitted: January 24, 2021; Peer review completed: May 8, 2021; Revisions accepted: May 12, 2021; Published: May 14, 2021.
DOI: 10.15376/biores.16.3.Bailon-Salas